

AD-A173 853

MODIFICATION OF CHEMICAL REACTIVITY VIA INCLUSION

1/1

COMPLEX FORMATION: PHOTO (U) COLUMBIA UNIV NEW YORK

DEPT OF CHEMISTRY S N RAO ET AL. 1986 AFOSR-TR-86-0993

UNCLASSIFIED

AFOSR-84-0040

F/G 7/3

NL


END

DATE

FORMED

1986

BY



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

Unclassified

MAR 31 1986

②

SECURITY CLASSIFICATION OF THIS PAGE

AD-A173 053

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified		1b. RESTRICTIVE MARKINGS										
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; Distribution Unlimited										
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S) <b>AFOSR-TR- 86-0993</b>										
6a. NAME OF PERFORMING ORGANIZATION Department of Chemistry Columbia University	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Office of Scientific Research										
6c. ADDRESS (City, State and ZIP Code) New York, New York 10027		7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling Air Force Base, D.C. 20332										
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Office of Sci. Res.	8b. OFFICE SYMBOL (If applicable) NC	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-84-0040										
8c. ADDRESS (City, State and ZIP Code) Bldg 410 Bolling Air Force Base, D.C. 20332		10. SOURCE OF FUNDING NOS. <table border="1"><tr><td>PROGRAM ELEMENT NO. 61102F</td><td>PROJECT NO. 2303</td><td>TASK NO. B2</td><td>WORK UNIT NO.</td></tr></table>		PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2303	TASK NO. B2	WORK UNIT NO.					
PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2303	TASK NO. B2	WORK UNIT NO.									
11. TITLE (Include Security Classification) Modification of Chemical Reactivity via Inclusion Complex Formation Photochemistry of <del>Dibenzyl Ketones and Benzyl Phenylacetates</del>												
12. PERSONAL AUTHOR(S) B.N. Rao, N.J. Turro, and V. Ramamurthy												
13a. TYPE OF REPORT reprint, scientific	13b. TIME COVERED FROM 1983 TO 85	14. DATE OF REPORT (Yr. Mo., Day) March, 1986	15. PAGE COUNT 4									
16. SUPPLEMENTARY NOTATION Journal of Organic Chemistry, vol. 51, 1986, pages 460-465												
17. COSATI CODES <table border="1"><tr><td>FIELD</td><td>GROUP</td><td>SUB. GR.</td></tr><tr><td></td><td></td><td></td></tr><tr><td></td><td></td><td></td></tr></table>		FIELD	GROUP	SUB. GR.							18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) dibenzyl ketone; benzyl phenylacetate; cyclodextrin; radical pairs; organic photochemistry; photolysis	
FIELD	GROUP	SUB. GR.										
19. ABSTRACT (Continue on reverse if necessary and identify by block number) In the context of employing inclusion complexes as a medium for organic photochemical reactions, we have investigated the photochemical behavior of dibenzyl ketones and benzyl phenylacetates using deoxycholic acid (DCA), Dianin's compound, and cyclodextrin (CD) as hosts in the solid state. Results on cage effect suggest that the translational motion of the benzyl radical pairs is restricted in all three media and totally in Dianin's compound. Products resulting from rearrangement of dibenzyl ketones were formed upon photolysis in Dianin's compound and cyclodextrin and were absent in deoxycholic acid. The absence of rearrangement in DCA and its presence to varied extents in Dianin's compound and cyclodextrin is suggested to be an indication of the restriction imposed by the host on the reorientational process of geminate radical pairs.												
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input checked="" type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION Unclassified										
22a. NAME OF RESPONSIBLE INDIVIDUAL Anthony Matuszko		22b. TELEPHONE NUMBER (Include Area Code) (202) 783-4963	22c. OFFICE SYMBOL NC									

DD FORM 1473, 83 APR

EDITION OF 1 JAN 73 IS OBSOLETE.

Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE

AIR FORCE OFFICE OF SCIENTIFIC AND TECHNICAL INFORMATION  
 AFOSR-TR-86-0993  
 This report was prepared by the  
 Department of Organic Chemistry, Indian Institute of Science, Bangalore 560012, India  
 and the Department of Chemistry, Columbia University, New York, New York 10027  
 under the sponsorship of the Air Force Office of Scientific and Technical Information Division

Accession For	
NTIS GRA&I	X
DTIC TAB	
Unannounced Justification	
By	
Distribution/	
Availability Codes	
Dist	Special
A-1 20	

## Modification of Chemical Reactivity via Inclusion Complex Formation: Photochemistry of Dibenzyl Ketones and Benzyl Phenylacetates

Bantu Nageswara Rao,<sup>†</sup> Nicholas J. Turro,<sup>\*†</sup> and Vaidhyanathan Ramamurthy<sup>\*†</sup>

Department of Organic Chemistry, Indian Institute of Science, Bangalore-560012, India, and Department of Chemistry, Columbia University, New York, New York 10027

Received August 2, 1985

In the context of employing "inclusion complexes" as a medium for organic photochemical reactions, we have investigated the photochemical behavior of dibenzyl ketones and benzyl phenylacetates using deoxycholic acid (DCA), Dianin's compound, and cyclodextrin (CD) as hosts in the solid state. Results on cage effect suggest that the translational motion of the benzyl radical pairs is restricted in all three media and totally in Dianin's compound. Products resulting from rearrangement of dibenzyl ketones were formed upon photolysis in Dianin's compound and cyclodextrin and were absent in deoxycholic acid. The absence of rearrangement in DCA and its presence to varied extents in Dianin's compound and cyclodextrin is suggested to be an indication of the restriction imposed by the host on the reorientational process of geminate radical pairs.

In the context of the study of reactivities of guest molecules and the way in which this reactivity is governed by the environment, the field of "inclusion complex" holds considerable promise. An essential characteristic of the host in the inclusion complex is its ability to form a solid structure with hollow spaces of large enough dimension to house prospective guests. Crystalline inclusion compounds may be subclassified as true clathrate-type, in which the guest molecules are imprisoned in discrete closed cavities or cages (Dianin's compound, cyclodextrin), the channel-type, in which the guest species are accommodated in continuous canals running through the crystal (deoxycholic acid, urea), and the layer-type where the guest component is situated between bands of the host structure (graphite). Zeolites belong to an intermediate class possessing cavities interconnected by channels. Significant changes in reaction product ratios have been effected by carrying out photochemical reactions in DCA and related matrices.<sup>1</sup> Thus the way was opened for employing in-

clusion complexes as a medium for organic photochemical studies.<sup>2</sup> Results described below on photochemical studies of dibenzyl ketones and benzyl phenylacetates using deoxycholic acid (DCA), Dianin's compound, and cyclodextrin (CD) as the host in the solid state represent our further interest in exploring the "host-guest" chemistry

- (1) Friedman, N.; Lahav, M.; Leiserowitz, L.; Popovitz-Biro, R.; Tang, C. P.; Zaretskii, Z. *J. Chem. Soc., Chem. Commun.* 1975, 864. Farina, M.; Audisio, G.; Natta, G. *J. Am. Chem. Soc.* 1967, 89, 5071. Brown, J. F.; White, D. M. *J. Am. Chem. Soc.* 1960, 82, 5671.
- (2) Turro, N. J.; Wan, P. *Tetrahedron Lett.* 1984, 25, 3655. Turro, N. J.; Wan, P. *J. Am. Chem. Soc.* 1985, 107, 678. Turro, N. J.; Cheng, C. C.; Lei, X. G.; Flanigen, E. M. *J. Am. Chem. Soc.* 1985, 107, 3739. Goswami, P. C.; de Mayo, P.; Ramnath, N.; Bernard, G.; Omkaram, N.; Scheffer, J. R.; Wong, Y. F. *Can. J. Chem.*, in Press. Padmanabhan, K.; Venkatesan, K.; Ramamurthy, V. *Can. J. Chem.* 1984, 62, 2025. Casal, H. L.; de Mayo, P.; Miranda, J. F.; Sciajno, J. C. *J. Am. Chem. Soc.* 1983, 105, 5155. Aoyama, H.; Miyazaki, K.; Sakamoto, M.; Omote, Y. *J. Chem. Soc., Chem. Commun.* 1983, 333. Tanaka, H.; Toda, F. *J. Chem. Soc., Chem. Commun.* 1983, 593. Popovitz-Biro, R.; Tang, C. P.; Chang, H. C.; Lahav, M.; Leiserowitz, L. *J. Am. Chem. Soc.* 1985, 107, 4043. Tang, C. P.; Chang, H. C.; Popovitz-Biro, R.; Frolow, F.; Lahav, M.; Leiserowitz, L.; McMullan, R. K. *J. Am. Chem. Soc.* 1985, 107, 4058. Arad-Yellin, R.; Brunie, S.; Green, B.; Knossow, M.; Tsoucaris, G. *J. Am. Chem. Soc.* 1979, 101, 7529.

<sup>†</sup> Indian Institute of Science.

<sup>†</sup> Columbia University.

Table I. Product Distribution upon Photolysis of Dibenzyl Ketones in Various Host Media in Solid State

host	guest:host <sup>a</sup> ratio	complex, mp, °C	% products <sup>b</sup>			RP <sup>c</sup>	cage effect, <sup>d</sup> %
			AA	AB	BB		
Dibenzyl Ketone							
α-CD	1:1.3		99			1	
β-CD	1:1	263	91			9	
γ-CD	1:1		81			19	
DCA	1:8	155	99			0	
Dianin's compd	1:45	169	95			5	
p-Methylbenzyl Benzyl Ketone							
β-CD	1:1.4	229	1	86	1.5	12	95
DCA	1:8	170	1.5	97	1.5	0	94
Dianin's compd	1:22	166-167		98		2	100
α-Methylbenzyl Benzyl Ketone							
β-CD	1:1	251	1.6	95	1.3	2	94
DCA	1:8	163	1.0	98	1.0	0	96
Dianin's compd	1:46	166-167		98		2	100

<sup>a</sup> Estimated on the basis of GC analysis; yields calculated with respect to reacted ketone. <sup>b</sup> Estimated by GC with an internal standard; error limit  $\pm 2\%$ . <sup>c</sup> RP corresponds to rearranged products 1-phenyl-4'-methylacetophenones and 1-phenyl-2'-methylacetophenones.

<sup>d</sup> Calculated according to the formula  $(AB - (AA + BB)) / (AA + AB + BB) \times 100\%$ .

Table II. Product Distribution upon Photolyses of Benzyl Phenylacetates in Various Host Media in Solid State

host	guest:host <sup>a</sup> ratio	complex mp, °C	% products <sup>b</sup>		
			AA	AB	BB
<i>p</i> -Methylbenzyl Phenylacetate					
β-CD	1:1	258	<1	≥99	<1
DCA	1:8	184	<1	≥99	<1
Dianin's compd	1:51	164	<1	≥99	<1
Benzyl ( <i>p</i> -Methoxyphenyl)acetate					
β-CD	1:1.3	262	<1	≥99	<1
DCA	1:8	185	<1	≥99	<1
Dianin's compd	1:51	166	<1	≥99	<1
<i>p</i> -Methylbenzyl ( <i>p</i> -Methoxyphenyl)acetate					
β-CD	1:1.4	256	<1	≥99	<1
DCA	1:8	182	<1	≥99	<1
Dianin's compd	1:54	163	<1	≥99	<1

<sup>a</sup> Estimated on the basis of GC analysis; yields calculated with respect to reacted ketone. <sup>b</sup> Estimated GC with an internal standard; error limit  $\pm 2\%$ .

to achieve selectivity in photochemical reactions.

The photodecarbonylation of dibenzyl ketones in homogeneous fluid solution occurs via a free-radical pathway in which coupling products (1,2-diarylethanes) are formed in quantitative yield.<sup>3</sup> For an unsymmetrical dibenzyl ketone (ACOB) the coupling products AA, AB, and BB are formed in the expected statistical ratio of 1:2:1. The solution photolysis of unsymmetrical benzyl phenylacetate yields 1,2-diarylethanes, but a marked cage effect has been noted in the more viscous solvents (e.g., 4-methylbenzyl 4-methoxyphenylacetate, 1:2.5:1 in dioxane and 1:5:1 in 2-propanol).<sup>4</sup> Photolysis of dibenzyl ketones and benzyl phenylacetates have been exploited extensively in several anisotropic environments—micellar solution,<sup>5</sup> silica surfaces,<sup>6</sup> uncoated glass surfaces,<sup>7</sup> and liquid crystalline media.<sup>8</sup> The reaction has been shown to result in enhanced cage recombination of geminate radical pairs and in marked <sup>13</sup>C enrichment of recovered starting ketones. Useful information regarding these anisotropic media has also been obtained through the use of the above reaction. In the present study the photochemical behavior of three dibenzyl ketones (1-3) and three benzyl phenyl acetates

(4-6) has been explored for understanding the influence of cavity or canal upon chemical reactions. Our results presented below indicate that generation of the two radicals A° and B° in the cavity or canal effectively sequesters these reactive species in which translational and rotational motions are restricted. This results in a dramatic enhancement of selectivity in formation of AB over AA and BB.

## Results

Addition of dibenzyl ketones and benzyl phenylacetates to saturated aqueous solutions of  $\beta$ -cyclodextrin precipitated a white solid. The solid was washed with excess of water and hexane and dried at 55 °C. Microcrystals of the inclusion complexes of DCA with the above carbonyl compounds were obtained by slow evaporation of methanol from an ice-cold solution containing excess of DCA. Dianin's compound complexes were prepared by slow crystallization from solutions of "empty" Dianin's compound in excess of liquid guest molecules. In some cases mild heating ( $\sim 40$  °C) was required to maintain the guest molecules in the molten state. DCA and Dianin's compound complexes were washed several times with excess of hexane and dried at 55 °C. All three complexes (cyclodextrin, DCA, and Dianin's compound) were obtained as white dry microcrystalline solids and were used as such for photolysis. Existence of true complexes in all cases was inferred from their mp, X-ray powder diffraction pattern, and solid-state NMR spectra. The X-ray powder pattern of the precipitated solid complex differed from those of

(3) Engel, P. J. *Am. Chem. Soc.* 1970, 92, 6074. Robbins, W. K.; Eastman, R. H. *J. Am. Chem. Soc.* 1970, 92, 6076.

(4) Givens, R. S.; Oettle, W. F. *J. Org. Chem.* 1972, 37, 4325.

(5) Turro, N. J.; Cherry, W. R. *J. Am. Chem. Soc.* 1978, 100, 7431.

(6) Frederick, B.; Johnston, L. J.; de Mayo, P.; Wong, S. K. *Can. J. Chem.* 1984, 62, 403.

(7) Baretz, B. H.; Turro, N. J. *J. Am. Chem. Soc.* 1983, 105, 1309.

(8) Hrovat, D. A.; Liu, J. H.; Turro, N. J.; Weiss, R. G. *J. Am. Chem. Soc.* 1984, 106, 5291.

the hosts. The solid-state  $^{13}\text{C}$  NMR spectra exhibited peaks corresponding to the guests in addition to the hosts. The host-guest ratio in each case was determined as follows. The amount of the guest present in a weighed sample of the complex was determined by GC using internal standard after extracting the guest with suitable solvent. Results are presented in Tables I and II. Close to a 1:1 ratio observed with cyclodextrin is consistent with our belief that the majority of the host molecules contain one guest molecule in their cavity. DCA is known<sup>9</sup> to form complexes with host-guest ratios ranging from 1:1 to 8:1. Formation of an 8:1 complex is known with long chain fatty acids ( $\text{C}_{15}\text{--C}_{29}$ ) as guests, and the canal length in these cases is believed to be  $\sim 28.9$  Å. Although, generally the host-guest ratio in the case of Dianin's compound<sup>10</sup> is  $\sim 6$ , the unexpectedly large value obtained here is an indication of either the poor method of complex preparation or distortion of the host structure by the guest molecule. It is quite likely that large size of the guests imposes strain on the system and thus keeps most of the pockets empty. Since none of the complexes gave suitable single crystals for X-ray crystallographic studies, no such investigation was pursued.

Photolyses of the solid complexes were done with 450-W medium-pressure mercury lamp either in degassed and sealed Pyrex tubes (in the case of dibenzyl ketones) or in quartz tubes (for benzyl phenylacetates). After irradiation, the products were extracted by dissolving the complex in suitable solvents and analyzed by GC. Product ratios in various media are summarized in Tables I and II. In addition to the decarbonylated products, small amounts of rearranged ketones 1-phenyl-4'-methylacetophenone and 1-phenyl-2'-methylacetophenone were obtained upon photolysis of dibenzyl ketone 1 in nonhomogeneous media such as micellar solution,<sup>11</sup> silica surface,<sup>6</sup> and zeolite.<sup>2</sup> Similar rearranged products are also expected for other dibenzyl ketones. In the present study, rearranged products were obtained in cyclodextrin and Dianin's compound and were absent (within GC detection limit, 1% of reacted ketone) in DCA matrix. Another interesting feature to come to light was the absence of cage escape products in Dianin's compound and their very minor amounts in cyclodextrin and DCA matrix ( $\sim 1\text{--}2\%$ ). However, with benzyl phenylacetates only one product (AB) was obtained in quantitative yield in all three media.

### Discussion

In the absence of structural data on the inclusion complexes under investigation we base our arguments on the structural details available for the inclusion complexes of DCA, Dianin's compound, and cyclodextrin with other guest molecules. The cavity/channel characteristics of these are briefly described below. DCA<sup>12</sup> crystallizes in such a manner as to leave a canal in which the guest can be closely confined. The DCA-guest interactions are hydrophobic, and thus the best fit depends principally on the size and shape of the guest molecule. The channel size in the *ab* plane is  $\sim 4 \times 6$  Å, and the length of the channel varies (*c* axis) depending on the guest molecule. The mole ratio of the DCA to guest component depends upon the size of the guest molecule and so varies from 1 to 8. Six molecules of Dianin's compound form a cage, and the cages are of





Figure 1. Cavity shapes and dimensions of the three host systems.



an hourglass shape that has been cut horizontally across the middle of each globe.<sup>13</sup> The cavity has a length of approximately 11 Å with a midpoint width of 4.2 Å and maximum upper and lower widths of 6.3 Å. The ends of each cage have a width of 2.8 Å and are formed by six hydrogen-bonded hydroxyl groups. Cyclodextrins,<sup>14</sup> one of the most commonly used "host" systems possess hydrophobic cavities that are able to include a variety of organic compounds whose character may vary from hydrophobic to ionic. Internal diameters and depth of cyclodextrins are:  $\alpha$ -cyclodextrin (4.2–8.8 and 7.8 Å),  $\beta$ -cyclodextrin (5.6–10.8 and 7.8 Å), and  $\gamma$ -cyclodextrin (6.8–12.0 and 7.8 Å) provide cavities for appropriately sized guest molecules. Cavity shapes and dimensions of the above three host systems that are used in the present study are illustrated in the Figure 1. While DCA and Dianin's compound belong to a category of polymolecular inclusion compounds, cyclodextrin belongs to monomolecular inclusion compounds.

Some of the interesting results that need elaboration are the following. The cage effect is total (100%) for benzyl phenylacetates, whereas it is so only in Dianin's compound for dibenzyl ketones. Very small amounts of noncage products ( $\sim 2\%$ ) are formed in the DCA and cyclodextrin matrixes (Table II). Rearranged products are formed only in Dianin's compound and cyclodextrin. Surprisingly they were absent in DCA medium. A mechanism for photolysis of 1–6 is presented in Scheme I. The difference in cage effect between benzyl phenylacetates and dibenzyl ketones may also be understood on the basis of the different spin states involved in the reaction. The decomposition of 1–3

(9) Herndon, W. C. *J. Chem. Educ.* 1967, 44, 724.

(10) Baker, W.; Floyd, A. J.; Meomie, J. F. W.; Pope, G.; Weaving, A. S.; Wild, J. H. *J. Chem. Soc.* 1956, 2010.

(11) Turro, N. J.; Chow, M. F.; Chung, C. J.; Weed, G. C.; Kraeutler, B. J. *Am. Chem. Soc.* 1980, 102, 4843.

(12) Jones, J. G.; Schwarzbaum, S.; Leisinger, L.; Low, B. W. *Acta Crystallogr., Sect. B* 1982, B38, 1207.

(13) Flippen, J. L.; Karle, J.; Karle, I. L. *J. Am. Chem. Soc.* 1970, 92, 3749.

(14) Saenger, W. *Angew. Chem., Int. Ed. Engl.* 1980, 19, 344.

is established<sup>3</sup> as being via triplet, while analogous esters<sup>4</sup> are believed to follow a singlet pathway. Long lifetime of the triplet benzyl radical pairs (derived from dibenzyl ketones) probably allows them to move farther in the host matrix and thus may result in a small percentage of non-cage products.

Unit cage effect observed for 1-3 in Dianin's compound is completely in accord with the structure of the Dianin's compound. Although the cavity size is large enough to accommodate 1-3, the ends of the cavity are narrow and therefore the radical pair cannot escape from the cavity. Indeed 100% cage effect is expected. Small noncage products (~2%) observed in DCA and cyclodextrin are unexpected as one would have expected 100% cage effect in solid matrix. However, the packing arrangements in both systems are such that there is a canal present. Therefore, a small percent of escape of the geminate radical pairs via the canals is not surprising. Another possibility is that a few percent of noncage products may also derive from small pockets of bulk liquid trapped by polycrystalline sample. This possibility cannot be ruled out on the basis of available structural information. Thus the results on cage effect suggest that the translational motion of the benzyl radical pairs is restricted in all three media.

Rearranged products (substituted acetophenone) were formed upon photolysis of dibenzyl ketones in Dianin's compound and cyclodextrin. These products have been proposed to arise via recombination of the original acyl benzyl radical pair with subsequent rearomatization (Scheme I).<sup>6,11</sup> In the case of dibenzyl ketone and  $\alpha$ -methylbenzyl benzyl ketone para and ortho coupling products are expected, whereas *p*-methylbenzyl benzyl ketone could give more than two, depending upon the site of cleavage. Only in the case of dibenzyl ketone was unequivocal identification of 1-phenyl-4'-methylacetophenone made by comparison with an authentic sample. In other cases products eluting in the GC very close to the starting ketone were presumed to be rearranged products. The opportunity for rearrangement would be enhanced by the longer lifetime of the geminate radical pair (phenylacetyl radical and benzyl radical) and by their freedom to undergo rotational motion within the restricted space. There is as yet little information about motional processes of guest molecule in complex clathrates. In the absence of such knowledge, chemical studies such as this can be informative. The large cage effect observed in all three media for 1-3 certainly suggests that the host medium is greatly restricting the movement of benzyl radicals. Therefore, the absence of rearrangement in DCA and its presence to varied extents in Dianin's compound and cyclodextrin is probably an indication of the restriction imposed by the host on the reorientational process of geminate radical pairs. The rotational process required for the rearrangement certainly seems to depend on the space available for the guest molecule. Yield of 1-phenyl-4'-methylacetophenone from dibenzyl ketone increases with the cavity size of the cyclodextrin ( $\gamma > \beta > \alpha$ ). Therefore absence of rearrangement in DCA is possibly due to the small size ( $4 \times 6$  Å) of the canal. Dianin's compound having slightly larger cavity (6.3-Å diameter) allows for rearrangement but not to the same extent as cyclodextrin (7-11-Å diameter).

Thus the results presented here illustrate that interesting effects can be obtained by carrying out photochemical reactions using organic host lattices. In view of the possibilities of predesigning a host from our knowledge of the thousands that have been proposed or prepared,<sup>15</sup>

it becomes immediately obvious that the field of selective photochemistry through clathration stretches limitlessly ahead.

## Experimental Section

**Materials.** Dibenzyl ketone (Aldrich) was twice distilled prior to use. Dibenzyl ketones 2 and 3 and benzyl phenylacetates 4-6 were prepared by following reported procedures.<sup>4,16</sup> All samples were purified by preparative TLC (silica gel, hexane-benzene). Cyclodextrins ( $\alpha$ ,  $\beta$ , and  $\gamma$ ) and deoxycholic acid were obtained from Sigma and were used as received. Dianin's compound was prepared by following the literature method.<sup>17</sup> Unsolvated Dianin's compound was obtained by dissolving the above compound in hot aqueous NaOH and passing CO<sub>2</sub> through the solution for about 30 min.

**Apparatus.** UV-vis absorption spectra were recorded on a Shimadzu UV-180 double-beam spectrometer. IR data were obtained with a Perkin-Elmer infrared spectrometer (Model 730). Varian T-60 and Bruker HW-270 FT NMR spectrometers were used for recording <sup>1</sup>H and <sup>13</sup>C NMR spectra, respectively. Solid-state <sup>13</sup>C NMR spectra were obtained by Dr. J. Ripmeester using a Bruker FT CXP-200 instrument. X-Ray powder photographs were recorded with a Phillips powder diffractometer employing monochromated Cu K $\alpha$  radiation. Gas chromatographic analyses were carried out on a Chemito GC instrument (Model 3800) fitted with a SE-30 5% column (8 ft  $\times$  1/8 in.) and a thermal conductivity detector. Irradiations were carried out with either a 450-W medium-pressure mercury lamp or a Rayonet reactor (RPR-100) fitted with 254-nm lamps.

**Preparation of Inclusion Complexes.** Cyclodextrin complexes were prepared as below. To a saturated aqueous solution of cyclodextrin, equimolar amounts of the guest molecule were added and stirred for 24 h. The white precipitate that formed was filtered, washed with cold water and ether, and dried at 50 °C for 10 h. Microcrystals of DCA complexes were obtained by slow evaporation of the solvent from a methanol solution of DCA (4 mM) and guest compound (1 mM). DCA complexes were washed several times with cold hexane and dried. Dianin's complexes with 1-6 were prepared by slow crystallization of unsolvated Dianin's compound from the guest liquids 1-6. In certain cases heating to 80 °C was required to dissolve the host. Microcrystalline complexes thus obtained were washed several times with hexane and dried. Inclusion complexes were characterized by their mp, X-ray powder photographs, and solid <sup>13</sup>C NMR spectra. The host-guest ratio was determined by GC analysis. The amount of 1-6 present in a known amount of the complex were determined by the GC integration of the guest peak with respect to the added internal standard (diphenyl ethane).

**Photolysis.** Irradiation of dibenzyl ketone complexes were carried out in a Pyrex tube using a 450-W medium-pressure mercury lamp and that of benzyl phenylacetates complexes in quartz tubes using a Rayonet reactor fitted with 254-nm lamps. Samples were degassed ( $10^{-3}$  mm) prior to irradiation and were rotated periodically during irradiation. Generally irradiations were stopped at ~20% conversion (8-12 h), and products were extracted as follows. Cyclodextrin complexes were dissolved in warm water and extracted with chloroform. DCA and Dianin's complexes were dissolved in hot aqueous methanol and extracted with hexane-chloroform mixture (9:1). Products were analyzed by GC and <sup>1</sup>H NMR.

**Acknowledgment.** V.R. thanks the Council of Scientific and Industrial Research, Government of India, and N.J.T. the National Science Foundation and the Air Force Office of Scientific Research for financial support. Dr. J. Ripmeester is thanked for <sup>13</sup>C NMR spectra of the solid complexes.

**Registry No.** Dibenzyl ketone-( $\alpha$ -CD) (1:1.3), 99766-03-7; dibenzyl ketone-( $\beta$ -CD) (1:1), 99765-87-4; dibenzyl ketone-( $\gamma$ -CD)

(15) Atwood, J. L.; Davies, J. E. D.; Mac Nicol, D. D., Eds. "Inclusion Compounds"; Academic Press: London, 1984; Vols. 1-3.

(16) Elderfield, R. C.; Burgess, K. L. *J. Am. Chem. Soc.* 1960, 82, 1975. Baratz, B. H.; Turro, N. J. *J. Am. Chem. Soc.* 1983, 105, 1304.

(17) Baker, W.; Floyd, A. J.; McOmie, J. F. W.; Pope, G.; Weaving, A. S.; Wild, J. H. *J. Chem. Soc.* 1956, 2010.

DATE  
FILMED  
2-8